

A Crossed Module giving the Godbillon-Vey Cocycle

Friedrich Wagemann

Institut Girard Desargues – ESA 5028 du CNRS

Université Claude Bernard Lyon-I

43, bd du 11 Novembre 1918

69622 Villeurbanne Cedex FRANCE

tel.: +33.4.72.43.11.90

fax: +33.4.72.43.00.35

e-mail: wagemann@desargues.univ-lyon1.fr

February 7, 2008

AMS classification: 17B56, 17B65, 17B66, 17B68

Abstract

We exhibit crossed modules (i.e. certain 4 term exact sequences) corresponding to the Godbillon-Vey generator for W_1 , $Vect(S^1)$ and $Vect_{1,0}(\Sigma)$.

Introduction

There is a well known correspondence in homological algebra between (equivalence classes of) exact sequences Λ -modules, starting in M and ending in N with n modules in between, and elements of $Ext_{\Lambda}^n(N, M)$. For $\Lambda = U(\mathfrak{g})$ the universal enveloping algebra of a Lie algebra $\mathfrak{g}$, this gives for example a correspondence between $H^2(\mathfrak{g}, V)$ and short exact sequences

$$0 \rightarrow V \rightarrow \hat{g} = V \times \mathfrak{g} \rightarrow \mathfrak{g} \rightarrow 0$$

where $\hat{g}$ is the semi-direct product of the abelian Lie algebra V and $\mathfrak{g}$. The Lie algebra law on $\hat{g}$ is given by a 2-cocycle of $\mathfrak{g}$ with values in V . Note that the short exact sequence is uniquely determined on the vector space level by V and $\mathfrak{g}$.

In the same way, there is a correspondence between $H^3(\mathfrak{g}, V)$ and certain 4 term exact sequences called crossed modules. In these sequences, only the first and the last term are specified, leaving (due to exactness) a choice of one $\mathfrak{g}$ -module to complete the sequence.

In this article, we exhibit crossed modules corresponding to the Godbillon-Vey 3-cocycle for W_1 , $Vect(S^1)$ and the 2-dimensional analogue of $Vect(S^1)$ $Vect_{0,1}(\Sigma)$ (or $Vect_{1,0}(\Sigma)$), Σ being a compact Riemann surface.

Acknowledgements

The author thanks J.-L. Loday for his question stimulating the present article and C. Kassel for making him aware of reference [6].

1 Crossed modules

In the same way as extensions, i.e. 3 term exact sequences, are related to Lie algebra cohomology in degree 2, crossed modules [6] [5], i.e. certain 4 term exact sequences, correspond to cohomology in degree 3.

Definition 1 *A crossed module of a Lie algebra is a homomorphism of Lie algebras $\mu : \mathfrak{m} \rightarrow \mathfrak{n}$ together with a $\mathfrak{n}$ -module structure η on $\mathfrak{m}$ such that*

- (a) $\mu(\eta(n) \cdot m) = [n, \mu(m)]$ for all $n \in \mathfrak{n}$ and all $m \in \mathfrak{m}$,
- (b) $\eta(\mu(m)) \cdot m' = [m, m']$ for all $m, m' \in \mathfrak{m}$.

One shows that $\ker(\mu) =: V$ is an abelian Lie algebra and that the action of $\mathfrak{n}$ on $\mathfrak{m}$ induces a structure of a $\mathfrak{g} := \text{coker}(\mu)$ -module on V . Note that in general $\mathfrak{m}$ and $\mathfrak{n}$ are not $\mathfrak{g}$ -modules.

An equivalence of crossed modules is defined in a natural way such that the restrictions of the maps on the kernel and the cokernel of μ are identical. Let us denote by $\text{crmod}(\mathfrak{g}, V)$ the set of equivalence classes of crossed modules. We have then the fundamental correspondence [6] theorem A.2, p. 138,

$$\text{crmod}(\mathfrak{g}, V) \cong H^3(\mathfrak{g}, V). \quad (1)$$

Let us briefly review how to associate to a crossed module a 3 cocycle of $\mathfrak{g}$ with values in V :

The exact sequence associated to a crossed module reads as follows:

$$0 \rightarrow V \xrightarrow{i} \mathfrak{m} \xrightarrow{\mu} \mathfrak{n} \xrightarrow{\pi} \mathfrak{g} \rightarrow 0$$

The first step is to take a (continuous) section σ of π and to calculate the defect of σ to be a Lie algebra homomorphism, i.e.

$$\alpha(x_1, x_2) := \sigma([x_1, x_2]) - [\sigma(x_1), \sigma(x_2)].$$

We have obviously $\pi \circ \alpha(x_1, x_2) = 0$, because π is a Lie algebra homomorphism, so $\alpha(x_1, x_2) \in \text{im}(\mu) = \ker(\pi)$. This means that there is a $\beta(x_1, x_2) \in \mathfrak{m}$ such that

$$\mu(\beta(x_1, x_2)) = \alpha(x_1, x_2).$$

Now, one can easily calculate that $\mu(d\beta(x_1, x_2, x_3)) = 0$ where d is the Lie algebra cohomology boundary of cohomology of $\mathfrak{g}$ with values in $\mathfrak{m}$ where $\mathfrak{g}$ acts on $\mathfrak{m}$ by $\eta \circ \sigma$.

This means that $d\beta(x_1, x_2, x_3) \in \ker(\mu) = \text{im}(i) = V$, i.e. there is a $\gamma(x_1, x_2, x_3) \in V$ such that $d\beta(x_1, x_2, x_3) = i(\gamma(x_1, x_2, x_3))$. It is fairly obvious that γ is a 3-cocycle of $\mathfrak{g}$ with values in V .

2 A crossed module giving $\theta \in H^3(W_1)$

In the following, we want to exhibit the 4 term exact sequence related to the Godbillon-Vey class in $H^3(W_1, \mathbb{R})$.

Recall that there is a semi-direct product of W_1 by its module of λ -densities. It is represented by a short exact sequence

$$0 \rightarrow F_\lambda \rightarrow F_\lambda \times W_1 \rightarrow W_1 \rightarrow 0,$$

where the Lie algebra structure on the product is given by

$$[(f, a), (g, b)] = ([f, g], L_f b - L_g a + c(f, g)).$$

Here, c is a 2-cocycle of W_1 with values in F_λ , the module of λ -densities on the line $\mathbb{R}$. We take $\lambda = 1$ and

$$c(f, g) = \begin{vmatrix} f' & g' \\ f'' & g'' \end{vmatrix}.$$

Now consider the short exact sequence of W_1 -modules

$$0 \rightarrow \mathbb{R} \rightarrow F_0 \xrightarrow{d} F_1 \rightarrow 0$$

Since W_1 acts by the Lie derivative $L_X = d \circ i_X + i_X \circ d$ on the density modules, the action commutes with the exterior differential d .

We can glue these 2 sequences together to get

$$0 \rightarrow \mathbb{R} \rightarrow F_0 \xrightarrow{d} F_1 \times W_1 \rightarrow W_1 \rightarrow 0, \quad (2)$$

d is trivially a Lie algebra homomorphism from the abelian Lie algebra F_0 to the semidirect product $F_1 \times W_1$. There is an action of $F_1 \times W_1$ on F_0 : it is given by the action of W_1 on F_0 . It gives a structure of a crossed module to the sequence (2). The second compatibility condition is trivial. The first one reads

$$d((f, a) \cdot b) = [(f, a), db]$$

The left hand side is just $d(L_f b) = d \circ i_{f \frac{d}{dt}} \circ db$. The right hand side gives $(0, L_f(db))$ and commutation of d and L_f shows condition (a). In conclusion, the 4 term exact sequence gives a crossed module. We claim that the crossed module corresponds to the Godbillon-Vey class via the isomorphism (1).

In fact, this is easy: one just has to move up the arrows in the wrong direction. First, one has to choose a section of the arrow $F_1 \times W_1 \rightarrow W_1$. But calculating the defect of this section to be a Lie algebra map just gives the cocycle of the semi-direct product. Then we have to move up the de Rham differential d , i.e. we have to choose a primitive. Afterwards, we have to take the Lie algebra coboundary. So, if we denote by a hat the primitive, we have to calculate

$$d(\hat{c})(f, g, h) = d\left(\widehat{\begin{vmatrix} -' & -' \\ -'' & -'' \end{vmatrix}}\right)(f, g, h).$$

But it is obvious that the Lie algebra coboundary commutes with taking the primitive. Furthermore, it is well known that we have the following relation

$$d\left(\begin{vmatrix} -' & -' \\ -'' & -'' \end{vmatrix}\right)(f, g, h) = \left(\begin{vmatrix} f & g & h \\ f' & g' & h' \\ f'' & g'' & h'' \end{vmatrix}\right)'$$

This expresses that fact that the Gelfand-Fuks cocycle is obtained from the Godbillon-Vey cocycle by “integration over the manifold” (and is easily checked directly).

Thus, we can choose the primitive such that the 3-cocycle associated to the above 4 term exact sequence is the Godbillon-Vey cocycle which is the generator of $H^3(W_1, \mathbb{R})$.

Remarks:

1) This reasoning also shows that $H^2(W_1, F_1)$ which is known to be 1-dimensional (consequence of Goncharova’s theorem, cf [2] p. 120), is generated by the cocycle

$$c(f, g) = \begin{vmatrix} f' & g' \\ f'' & g'' \end{vmatrix} dx.$$

2) We owe to C. Roger the remark that the crossed module is obtained under Yoneda product $Ext_{U(W_1)}^1(F_1, \mathbb{R}) \times Ext_{U(W_1)}^1(\mathbb{R}, F_1) \rightarrow Ext_{U(W_1)}^2(\mathbb{R}, \mathbb{R}) \cong H^3(W_1, \mathbb{R})$. This possibility bears perhaps deep homological properties.

3 A crossed module giving $\theta \in H^3(Vect(S^1), \mathbb{C})$

Now we want to generalize the preceeding situation to the circle. There is an obvious problem to do so: the sequence

$$0 \rightarrow \mathbb{C} \rightarrow \mathcal{F}_0 \xrightarrow{d} \mathcal{F}_1 \rightarrow 0$$

where $\mathcal{F}_0 = \mathcal{C}^\infty(S^1)$ and $\mathcal{F}_1 = \Omega^1(S^1)$ is not exact, but has a cokernel $H^1(S^1, \mathbb{C}) = \mathbb{C}$.

In order to get around this problem, let us recall some evident, but somewhat strange facts:

There is a chain of inclusions of Lie algebras

$$W_1^{pol} \hookrightarrow Vect(S^1) \hookrightarrow Vect(\mathbb{R})$$

Here, $W_1^{pol} = \bigoplus_{n \geq -1} \mathbb{C}x^{n+1} \frac{d}{dx}$ is the Lie algebra of (complexified) polynomial vector fields on the line, $Vect(S^1) = \widehat{\bigoplus_{n \in \mathbb{Z}} \mathbb{C}e^{int} \frac{d}{dt}}$ is the Lie algebra of complexified vector fields on the circle (the hat meaning that $e^{int} \frac{d}{dt}$ for $n \in \mathbb{Z}$ is a topological basis, because periodic functions can be approximated by Fourier polynomials), and $Vect(\mathbb{R}) = \{f(x) \frac{d}{dx} \mid f \in \mathcal{C}^\infty(\mathbb{R}, \mathbb{C})\}$ is the Lie algebra of complexified vector fields on the real line $\mathbb{R}$.

The maps $x^{n+1} \frac{d}{dx} \mapsto -ie^{int} \frac{d}{dt}$ (i.e. one sets $x = e^{it}$) and $f(t) \frac{d}{dt} \mapsto \tilde{f}(x) \frac{d}{dx}$ (where the field $f(t) \frac{d}{dt}$ is lifted to the universal covering to the unique 1-periodic field $\tilde{f}(x) \frac{d}{dx}$) are easily seen to be Lie algebra homomorphisms.

The strange fact is that W_1^{pol} and $Vect(\mathbb{R})$ have the same continuous cohomology, but $Vect(S^1)$ does not. This is elucidated by the fact that the isomorphism in cohomology between W_1^{pol} and $Vect(\mathbb{R})$ is not induced by the above inclusion, but by the “Taylor expansion at 0”-map $Vect(\mathbb{R}) \rightarrow W_1^{pol}$ (which is a continuous surjection of Fréchet spaces by Borel’s lemma). One sees that W_1^{pol}

and $Vect(\mathbb{R})$ correspond to the 2 ways of associating $\mathbb{R}$ to S^1 : as its tangent space, or as its universal covering.

I recall all this just to motivate the fact that $Vect(S^1)$ acts on F_λ (the λ -densities on $\mathbb{R}$) and W_1^{pol} acts on $\mathcal{F}_\lambda$ (the λ -densities on S^1). In the first case, we embed $Vect(S^1)$ into $Vect(\mathbb{R})$, and in the second case, we embed W_1^{pol} into $Vect(S^1)$.

This gives us the possibility to consider the exact sequence

$$0 \rightarrow \mathbb{R} \rightarrow F_0 \xrightarrow{d} F_1 \times Vect(S^1) \rightarrow Vect(S^1) \rightarrow 0.$$

By the same arguments as above, this is a crossed module giving the Godbillon-Vey cocycle.

4 A crossed module giving $\theta \in H^3(Vect_{1,0}(\Sigma), \mathbb{C})$

Now, let us recall that $H^3(Vect_{1,0}(\Sigma), \mathbb{C})$, where $Vect_{1,0}(\Sigma)$ is the Lie algebra of differentiable vector fields of type $(1, 0)$ (i.e. locally, such a field is written $f(z, \bar{z}) \frac{\partial}{\partial z}$) on a compact Riemann surface Σ , is also of dimension 1 and generated by a Godbillon-Vey type cocycle, see [1], [8]. This cocycle is the integral over Σ over a determinant as in the usual Godbillon-Vey cocycle, all derivatives being with respect to z .

We propose a crossed module corresponding to this generator.

There are two obvious problems: first of all, there is the same problem as in the preceding section, so we have to lift our fields to the universal covering, and second, there is the problem that the derivatives involved are only ∂ , instead of being $d = \partial + \bar{\partial}$. For the second problem, we define a new action of $Vect_{1,0}(\Sigma)$ on $\bigoplus_{i=0,1} \Omega^{i,0}(\Sigma)$ by the modified Cartan formula:

$$L_X \omega = i_X \circ \partial + \partial \circ i_X.$$

This means explicitly

$$L_X \omega = \begin{cases} i_X(\partial(\omega)) & \text{if } \deg(\omega) = 0 \\ \partial(i_X(\omega)) & \text{if } \deg(\omega) = 1 \end{cases}$$

One calculates easily that this defines a Lie algebra action of $Vect_{1,0}(\Sigma)$ on $\bigoplus_{i=0,1} \Omega^{i,0}(\Sigma)$ and that

$$0 \rightarrow \mathbb{C} \rightarrow \Omega^{0,0}(\Sigma) \xrightarrow{\partial} \Omega^{1,0}(\Sigma)$$

is an exact sequence of $Vect_{1,0}(\Sigma)$ -modules, unfortunately the last map is not surjective. Now, let us lift these modules to the universal covering:

Suppose $g > 0$ in order to have as the universal covering of Σ either $\mathbb{C}$ or $\mathbb{H}$, and let U denote $\mathbb{C}$ in case the genus is 1 and $\mathbb{H}$ in case the genus is greater than 1. We have a short exact sequence of $Vect_{1,0}(\Sigma)$ -modules

$$0 \rightarrow \mathbb{C} \rightarrow \Omega^{0,0}(U) \xrightarrow{\partial} \Omega^{1,0}(U) \rightarrow 0$$

Now we can form the 4 term exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \Omega^{0,0}(U) \xrightarrow{\partial} \Omega^{1,0}(U) \times Vect_{1,0}(\Sigma) \rightarrow Vect_{1,0}(\Sigma) \rightarrow 0$$

This sequence gives obviously by the same arguments as before a crossed module with corresponding 3-cohomology class represented by the Godbillon-Vey cocycle.

Note that the same construction works for the Lie algebra of holomorphic vector fields on an open Riemann surface $Hol(\Sigma_r)$ where $\Sigma_r = \Sigma \setminus \{p_1, \dots, p_r\}$ acting on the module of holomorphic λ -densities $\mathcal{F}_\lambda(\Sigma_r)$ by the modified Cartan formula. This gives a crossed module describing $H^3(Hol(\Sigma_r), \mathbb{C})$ which is also known to be 1-dimensional [4].

References

- [1] Feigin, B. L.: Conformal field theory and Cohomologies of the Lie algebra of holomorphic vector fields on a complex curve. Proc. ICM, Kyoto, Japan, 71-85 (1990)
- [2] Fuks, D. B.: Cohomology of Infinite Dimensional Lie algebras. New York and London: Consultant Bureau 1986
- [3] Gelfand, I. M., Fuks, D. B.: Cohomologies of Lie Algebra of Tangential Vector Fields of a Smooth Manifold. Funct. Ana. and its Appl. **3** (1969) 194-210
- [4] Kawazumi, N.: On the Complex Analytic Gel'fand-Fuks Cohomology of Open Riemann Surfaces. Ann. Inst. Fourier, Grenoble **43** 3 (1993) 655-712
- [5] Loday, J.-L.: Cyclic Homology. Springer Grundlehren **301** 1991
- [6] Loday, J.-L., Kassel, C.: Extensions centrales d'algèbres de Lie. Ann. Inst. Fourier, Grenoble **32**, 4 (1982) 119-142
- [7] Wagemann, F.: Differential Graded Cohomology and Lie Algebras of Holomorphic Vector Fields. Comm. Math. Phys. **208** (1999) 521-540
- [8] Wagemann, F.: A Two Dimensional Analogue of the Virasoro Algebra. to appear in J. Geo. Phys.